



International Journal of Pharmaceutics and Drug Analysis

Content Available at www.ijpda.com

ISSN: 2348:8948



MICROEMULSIONS: A PROMISING DRUG DELIVERY SYSTEM

Adarsh Kumar, Kapil Kumar, Aparna Joshi, Vaishali Rajput

Division of Pharmaceutics, Global Institute of Pharmaceutical Education and Research, Kashipur, Uttarakhand, India.

Received: 04 Sept 2023 Revised: 25 Sept 2023 Accepted: 05 Nov 2023

Abstract

Microemulsions, comprising oil, water and a surfactant, in association with some co-surfactant, are thermodynamically stable systems. They have found applications in a large number of chemical and pharmacological processes due to their unique properties such as large interfacial area, low interfacial tension, and most importantly, the ability to solubilize and deliver hydrophobic drugs. Microemulsions are oral, topical and parenteral administration. Microemulsions are thermodynamically stable. Microemulsions are clear, thermodynamically stable isotropic liquid mixtures of oil, water and surfactant, frequently in combination with a cosurfactant. This review highlights the properties and several recent developments in the use of microemulsions for medical treatment purposes including targeted drug delivery.

Keywords: Co-surfactants, Microemulsion, Surfactants, Thermodynamically stable.

This article is licensed under a Creative Commons Attribution-Non-commercial 4.0 International License. Copyright © 2023 Author(s) retains the copyright of this article.



*Corresponding Author

Vaishali Rajput

DOI: <https://doi.org/10.47957/ijpda.v11i4.562>

Produced and Published by
[South Asian Academic Publications](http://SouthAsianAcademicPublications.com)

Introduction

Microemulsions are merely alike emulsions that are a different class of colloidal systems. We can observe colloidal systems in all three possible states of matter—gas, liquid, and solid. Basically, the emulsions are formed via the emulsification of two or more immiscible liquids primarily.

The emulsions are basically formed from two or more immiscible liquid phases via emulsification, a profound characteristic of surface-active agents, making them versatile for a wide range of practical application, for instance, milk and cream, butter, margarine, mayonnaise, espresso, cutting fluid for metal working, and the photo-sensitizer. The two types of microemulsions are direct (oil dispersed in water o/w) & reversed (water dispersed in oil, w/o) [1-5].

Various techniques have been published in the scientific literature to enhance the dissolution profile and also the absorption efficiency and bioavailability of water-insoluble and liquid lipophilic drugs [6-10].

- Reduction of partical size via micro-nization.
- Co-grinding

- Adsorption on porous structures
- Solid dispersion
- Self-emulsification
- Microencapsulation

1. Structure of Microemulsion




Figure 1: Microemulsion Structure

Microemulsions or Micellar emulsion are dynamic system in which the interface is continuously and spontaneously fluctuating [11-13]. Structurally, they are divided in to oil in water (o/w), water in oil (w/o) and bi-continuous microemulsions. In w/o microemulsions, water droplets are dispersed in the continuous oil phase while o/w microemulsions are formed when oil droplets are dispersed in the continuous aqueous phase. In system where the amounts of water and oil are similar, the bi-continuous microemulsions may result [14-18]. The mixture oil water and surfactants are able to form a wide variety of structure and phase depending upon the proportions of component.

2. Advantages of Microemulsion Based Systems [19,20]

Microemulsions exhibit several advantages as a drug delivery system:

- i. Microemulsions are thermodynamically stable systems and allows self-mediations of this system.
- ii. Microemulsions are both solubilised in hydrophilic and lipophilic drugs.
- iii. The dispersed phase, lipophilic or hydrophilic (oil in water, o/w, or water in oil, w/o microemulsions) can act as a potential reservoir of hydrophilic or lipophilic drugs.
- iv. Microemulsions have low viscosity compared to, primary and multiple emulsions.

3. Disadvantages of Microemulsions Based Systems: [21,22]

- i. Require large amount of S/Cs for stabilizing droplets.
- ii. The surfactant should be nontoxic for use in pharmaceutical applications.
- iii. Microemulsion stability is influenced by environmental parameters such as temperature and pH.

4. Limitations: [23,24]

Factors which limit the use of microemulsion in pharmaceutical applications:

- i. The concentration of surfactants and co-surfactants used must be kept low for toxicological reasons.
- ii. Microemulsion also suffers from limitations of phase separation.
- iii. For intravenous use, the demand of toxicity on the formulation is rigorous and very few studies have been reported so far.

Depending upon the nature of the dispersed particles, microemulsions are classified into the following types:

1) Water-in-oil microemulsion

The W/O type microemulsion is a dispersion of water or an aqueous solution in a water-immiscible liquid. The water is, in this case, the "discontinuous" (inner) phase, and the oil is the "continuous" (outer) phase (Figure 2).

2) Oil-in-water microemulsion

he oil-in-water type is a dispersion of a water-immiscible liquid (always called the oil) in an aqueous phase. The oil is, in this case, the "discontinuous" (inner) phase, and the aqueous phase is the "continuous" (outer) phase (Figure 2).

3) Bicontinuous microemulsion

It consists of an infinite bilayer bent everywhere with a saddle-like curvature so that it is multiconnected to itself (by many random "passages" similar to tubular connections in the sponge) isotropically over macroscopic distances and divides space into two independent solvents. Bicontinuous structures are present in the systems where the amount of water and oil is similar. Both the oil and water exist as a continuous phase in the presence of a continuously fluctuating surfactant-stabilized interface with a net curvature of zero

Methods of preparation of Microemulsion [27,28]

Two different methods are utilized to prepared microemulsion.

7.1. Phase Titration Method:

Microemulsions are prepared by the spontaneous emulsification method (phase titration method). Construction of phase diagram is a useful approach to study the complex series of interactions that can occur when different components are mixed. Microemulsions are formed along with various association structures (including emulsion, micelles, lamellar, hexagonal, cubic, and various gels and oily dispersion) depend on the chemical composition and concentration of each component.) The region can be separated into w/o or o/w microemulsion by simply considering the composition it is oil rich or water rich. [29]

7.2.2 Phase Inversion Method:

Phase inversion of microemulsion method, physical changes can occur, also changes in particle size, these can be ultimately affected drug release in in-vitro and in-vivo for non ionic surfactants can be accomplish by the changing the temperature of the system, in these processes an o/w microemulsion at low temperature changes to w/o microemulsion. This is also called as transitional phase inversion method. During the cooling, the system cross the zero-point spontaneous shape and maintaining the surface tension, and increasing the formation of oil droplet dispersion[30].

In this phase inversion method, transition in the radius can be occur by changing in the water volume fraction. Water volume fraction can be increased, surfactants from stabilizing a w/o microemulsion to an o/w microemulsion using temperature.

Evaluation of Microemulsion [31,32,33]

- Visual Inspection
- Thermodynamic stability
- Measurement of p^H
- Viscosity measurements
- Zeta potential determination
- Particle size determination
- Drug content estimation

- **Visual Inspection**

Visual inspection was made after each addition of water to the oil and surfactant and co-surfactant mixture. The samples were identified as microemulsion, emulsion or gel formation by visual observation.

- **Thermodynamic stability**

To overcome the problem of metastable formulation, thermo-dynamic stability tests were performed.

a) Centrifugation

The formulation was centrifuged at 3500 rpm for 30 min to ensure physical stability.

b)Stress test

These tests were done to optimize the best microemulsion formulation under extreme conditions. Stress was carried out at 4 °C and 45 °C for 48 h each for a period of six cycles, followed by 25 °C and 21 °C for 48 h for about three cycles. The samples were checked for coalescence, cracking or phase separation.

- **Measurement of pH**

The pH values of the optimized formulation were measured by immersing the electrode directly into the dispersion using a calibrated pH meter (Digital Potentiometer Model EQ-601 Equip-Tronics).

- **Viscosity measurements**

The viscosity of the optimized formulation was determined as such without dilution using Brookfield Viscometer (DV-E Brookfield Viscometer Model-LVDVE).

- **Zeta potential determination**

Zeta potential of samples was measured by Zeta sizer. Samples were placed in clear disposable zeta cells and results were recorded. Before putting the fresh sample, cuvettes were washed with methanol and rinsed using the sample to be measured before each experiment.

- **Particle size determination**

The mean particle size and particle size distribution of drug loaded microemulsion were determined by Horiba SZ-100 nanoparticle analyzer, at 28°C. It measures the fluctuation of the intensity of the scattered light which is caused by particle movement. Each sample was measured in triplicate.

- **Drug content estimation**

Microemulsion containing 100 mg drug was dissolved in 100 ml 0.1N HCl taken in volumetric flask. Then the solvent was filtered, 1 ml was taken in 50 ml volumetric solution and diluted up to the mark with 0.1N HCl and analyzed spectrometrically at 295 nm. The concentration of drug in mg/ml was obtained by using standard calibration curve of the drug. Drug content studies were carried out in triplicate for each formulation batch.

5. Marketed formulation of Microemulsion:

Ascabiol emulsion, Noreva Sebodiane DS, Winsor ointment etc.

6. Application of Microemulsion:

These are some applications of microemulsions in delivery of drug.

During the last two decades, microemulsions are used as drug delivery system, they are offer the advantages like thermodynamic stability, optical clarity, easy of penetration

Parenteral Delivery

The formulation of parenteral dosage form of lipophilic

and hydrophilic drugs has proven to be difficult. The formulation of w/o microemulsions are beneficial in the parenteral delivery of sparingly soluble drugs where the administration of suspensions is not required. For frequent administration of drugs requires high concentration. They are existing the higher physical stability in plasma than liposomes or other vehicles and the internal oil phase is more resistant against drug leaching. Several sparingly soluble drugs have been formulated in to o/w microemulsions for parenteral delivery. Von Corsewant and Thoren was taken the alternate approach in which C3- C4 alcohols were replaced with parenterally acceptable co- surfactants, poly ethylene glycol (400)/ poly ethanol glycol (600) 12-hydroxy stearate/ ethanol will maintaining a flexible surfactant film, these are forming the spontaneous curvature near zero, in the microemulsions obtaining the almost balanced middle phase [34].

Oral Delivery

The development of effective oral delivery systems has been challenging because drug efficiency can be restricted by instability or poor solubility in gastrointestinal fluid. The solubilization of poorly soluble drugs (particularly BCS class II/class IV) are enhanced by the microemulsions. Also overcome the dissolution related bioavailability problems. The presence of polar, non-polar and interfacial domains, hydrophilic drugs are encapsulated with varying solubility of macromolecules. These systems have been protecting the incorporated drugs agonist oxidation, enzymatic degradation and enhance membrane permeability. Commercially available microemulsions for formulation of oral delivery are Sandimmune Neoral® (Cyclosporine A), Fortovase® (Saquinavir), Norvir® (Ritonavir) etc. Improving the oral bioavailability of poorly water-soluble drugs can be enhanced their solubility in gastrointestinal fluids are potential useful in the formulation of microemulsions [35].

Topical Delivery

Topical administration of drugs having the advantages like avoidance of hepatic first-pass metabolism of drug and related toxicity effects. These are having the direct delivery and targetability of the drug to affected areas of the skin and eyes. The area of drug penetration into the skin having the number of studies. In these drug penetration studies they are incorporate both hydrophilic (5-fluorouracil, apomorphine hydrochloride, diphenhydramine hydrochloride, tetracaine hydrochloride, methotrexate) and lipophilic drugs (oestradiol, finasteride, ketoprofen, meloxicam, felodipine, triptolide) and enhance their penetration. In the formulation of microemulsions requires high surfactant concentration. Skin irritating aspects must be considered specially when they are intended to be applied for a long period [36].

Nasal Delivery

Recently, microemulsions are being studied as a delivery system to enhance uptake of drug through nasal mucosa. Mucoadhesive polymers helps in prolonging residence time on the mucosa. Lianly et al. investigating the effect of diazepam on the emergency treatment of status epilepticus. They found that the nasal absorption of diazepam fairly rapid at 2 mg kg⁻¹ dose with maximum drug plasma concentration reached within 2-3 min [37].

Current and Future Developments

During the last two decades lot of research work has been carried out on microemulsion system (Table 1, 2) for providing novel solutions to overcome the problems of poor aqueous solubility of highly lipophilic drug compounds and provide reproducible bioavailability. Industrial point of view, it can be easily scaled up with considering relative cost of commercial production. Microemulsion can also be used for cosmetic purpose and drug targeting. Now a day, researcher work is focused on the production of safe, efficient and more compatible microemulsion constituents which will further enhance the utility of this novel delivery system [38-40].

Conclusion:

Microemulsions are having a vast and significant potential in drug delivery as well as in the industrial process. Researchers are working in this field for drug release, coatings, dyes, agrochemicals and in enzyme reaction. In the future prospects, microemulsions will be used in synthesis of nanoparticles and as a industrial chemical sensors. The role of microemulsion in providing novel solutions to overcome the problems of poor aqueous solubility of highly lipophilic drug compounds and provide high, more consistent and reproducible bioavailability. Furthermore, these formulations can be easily manufactured in term of the relative cost of commercial production.

Acknowledgments

Authors are thankful for the management of the institute to provide necessary facilities for this work.

Disclosure of conflict of interest

No conflict of interest is associated.

References

1. Shah Rohit Ramesh, Magdum Chandrakant Shripal; Preparation and Evaluation of Aceclofenac Topical Microemulsion; Iranian Journal of Pharmaceutical Research., 9(1) (2010) 5-11.
2. Sharma Pankaj Kumar; Enhancement of solubility and stability of Celecoxib using microemulsion based topical formulation; Journal of Pharmacy Research., 4(7) (2011) 2216-2220.
3. Jadhav K. R, Shetye S. L, Kadam V. J; Design and Evaluation of Microemulsion Based Drug Delivery

- System; International Journal of Advances in Pharmaceutical Sciences., 1 (2010) 156-166.
4. Fathy I. Abdallah, Hamdy M. Dawaba; Evaluation of the anti-inflammatory and analgesic effects of piroxicam loaded microemulsion in topical formulations; Int J Pharm Pharm Sci., 3(2)(2011) 6670
5. Shishu, Rajan Sunita and Kamalpreet; Development of Novel Microemulsion-Based Topical Formulations of Acyclovir for the Treatment of Cutaneous Herpetic Infections; AAPS PharmSciTech., 10 (2009) 559-565.
6. Ikechukwu UR, John Francis DE, Ambi AA. Development and evaluation of Ritonavir hollow microballoons for floating drug delivery. Universal Journal of Pharmaceutical Research 2017; 2(2): 8-11.
7. Angela Attar Nasser, Reza Aboofazeli, Hossein Zia, Thomas E Needham; Lecithin Stabilized Microemulsion - Based Organogels for Topical Application of Ketorolac Tromethamine. II. In vitro Release Study; Iranian Journal of Pharmaceutical Research; (2003) 117-123.
8. Mrunali R. Patel, Rashmin B. Patel, Jolly R. Parikh; Effect of Formulation Components on the In Vitro Permeation of Microemulsion Drug Delivery System of Fluconazole; AAPS PharmSciTech., 10 (2009) 917-923.
9. Yusuf FS. Formulation and *in-vitro* evaluation of floating microballoons of stavudine. Universal Journal of Pharmaceutical Research 2016; 1(1): 8-11.
10. Susijit Sahoo, Bhakti Bhusan Barik, Nihar Ranjan Pani; Formulation and characterization of microemulsion of an anti-fungal drug; Journal of Pharmacy Research., 4(7) (2011) 2397-2399.
11. Nadeem Abbas Kizilbash, Dhafallah Alenizi; Design of a Microemulsion-Based Drug Delivery System for Diclofenac Sodium; J.Chem.Soc.Pak., 33 (2011), 1-6.
12. Yuan Y, Li S-M, Mo F-K, D-F Zhong; Investigation of microemulsion system for transdermal delivery of meloxicam. Int J Pharm 2006; 321: 117-123.
13. Hellweg T, Phase structure of microemulsions, Curr opin colloid interface sci., 7, (2002) 50-56.
14. Kumar. K. Senthil, Dhachinamoorthi. D, Saravanan. R; Microemulsions as Carrier for Novel Drug Delivery: A Review; International Journal of Pharmaceutical Sciences Review and Research., 10 (2011) 37-45.
15. Chauhan N, Kumar K, Pant NC. An updated review on transfersomes: a novel vesicular system for transdermal drug delivery. Universal Journal of Pharmaceutical Research 2017; 2(4): 42-45.
16. Hunter, R. J; Introduction to Modern Colloid Science; 1st ed.; Oxford University Press: Oxford, 1994.
17. Sushama Talegaonkar, Adnan Azeem, Farhan Ahmad J, Roop Khar K, Shadab Pathan A, Zeenat Khan I; Microemulsions: A Novel approach to enhanced drug delivery; Recent patents on drug delivery and formulation; 2, 2008; 238-257.
18. Shafiq un Nabi S, Shakeel F, Talegaonkar S;

- Formulation development and optimization using nanoemulsion technique: A technical note; *AAPS Pharm Sci Tech*, 8 (2007) 1-6.
19. ALGIN YAPAR E, BESKAN U, KARAVANA SY. A recent overview of locally administered topical otic dosage forms. *Universal Journal of Pharmaceutical Research* 2019; 4(4): 39-42.
 20. Hsiu-O Ho, Chih-Chuan Hsiao, Ming-Thau Sheu, preparation of microemulsions using polyglycerol fatty acid esters as surfactant for the delivery of protein drugs. *J. pharm. Sci.* 1996; 58: 138-143.
 21. Dreher F, Walde P, Walther P, Wehrli E, Interaction of a lecithin microemulsion gel with human stratum corneum and its effect on transdermal transport. *J. Control. Rel.* 1997; 45:131-140.
 22. Agarwal P, Semimul A. A comprehensive review on sustained release matrix tablets: a promising dosage form. *Universal Journal of Pharmaceutical Research* 2018; 3(6): 49-54.
 23. Ashok Patel R, Pradeep vavia R; Preparation and In vivo Evaluation of Self-Microemulsifying Drug Delivery System Containing fenofibrate; *The AAPS Journal*, 9 (2007) 344-352.
 24. Peltola S, Saarinen S P, Kiesavaara J ,Urttia S T M; Microemulsions for topical delivery of estradiol; *Int J Pharm.*, 254 (2003) 99-107.
 25. Alfisha Saifi, Kapil Kumar, Deepak Teotia. A basic review on micro emulsion as a drug delivery system. *Journal of Biomedical and Pharmaceutical Research*. Volume 9, Issue 5: September-October: 2020, 06-11.
 26. Rahi N, Kumar K. Nanosponges: a new era of versatile drug delivery system. *Universal Journal of Pharmaceutical Research* 2017; 2(3): 30-33.
 28. 2Gurleen Kaur, Alfisha Saifi, Kapil Kumar , Deepak Teotia. Development and Evaluation of Micro Emulsion Formulations of Nebivolol for Solubility Enhancement, *Journal of Drug Delivery and Therapeutics*. 2021; 11(5):84-89.
 29. Osama Umar, Kapil Kumar, Aparna Joshi, Dipti Khairiya, Deepak Teotia, Ikram. A comprehensive review on microemulsions: a potential novel drug delivery system. *Int J Indig Herbs Drugs* 2022; 7(3): 56-61.
 30. Fialho SL, da Silva-Cunha A. New vehicle based on a microemulsion for topical ocular administration of dexamethasone. *Cl i n Ex p e r Op h t h a l m o l*, 2004; 32(6):626-632
 31. Syamasri Gupta, S.P. Moulik, Biocompatible microemulsions an their prospective uses in drug delivery. *Journal of Pharmaceutical Sciences*. 2008; 97:22-45.
 32. Yogeshwar G. Bachhav and Vandana B. Patravale; Microemulsion-Based Vaginal Gel of Clotrimazole: Formulation, In Vitro Evaluation, and Stability Studies; *AAPS PharmSciTech*; Vol. 10, No. 2, June 2009, 476-481.
 33. Peter OI, Ifeoma UC. Development and evaluation of Albendazole microcapsule for colonic drug delivery system. *Universal Journal of Pharmaceutical Research* 2017; 2(2): 4-7.
 34. E Gundogdu, I Gonzalez Alvarez, E Karasulu; Improvement of effect of water-in-oil microemulsion as an oral delivery system for fexofenadine: in vitro and in vivo studies; *International Journal of Nanomedicine*; 2011:6 1631-1640.
 35. Nweje-Anyalowu Paul C, Anyalogbu Ernest AA, White Alalibo Jim. Design and evaluation of chronotherapeutic pulsatile drug delivery system of Cilnidipine. *Universal Journal of Pharmaceutical Research* 2017; 2(5): 15-18.
 36. Amit A. Kale and Vandana B. Patravale; Development and Evaluation of Lorazepam Microemulsions for Parenteral Delivery; *AAPS PharmSciTech*,9 (2008) 966-971.
 37. Vandana Patel, Hirenkumar Kukadiya, Rajshree Mashru; Development of Microemulsion for Solubility Enhancement of Clopidogrel; *Iranian Journal of Pharmaceutical Research*, 9(4) (2010), 327-334.
 38. Kaur G, Paliwal S. Formulation and evaluation of etoricoxib microbeads for sustained drug delivery. *Universal Journal of Pharmaceutical Research* 2019; 4(1): 35-39.
 39. Park K M, Kim C K; Preparation and evaluation of flurbiprofen- loaded Microemulsions for parental delivery; *Int J Pharm.*, 181 (1999) 173-179.
 40. Sunday OS. Colon-targeted drug delivery systems: design, trends and approaches. *Universal Journal of Pharmaceutical Research* 2017; 2(4): 46-50.